

## Sea Turtle Abundance at Isolated Reefs of the Mariana Archipelago

S. P. KOLINSKI

Joint Institute for Marine and Atmospheric Research  
National Marine Fisheries Service, Pacific Islands Regional Office  
1601 Kapiolani Blvd., Suite 1110, Honolulu, HI 96814-0047

R. K. HOEKE, S. R. HOLZWARH AND P. S. VROOM

Joint Institute for Marine and Atmospheric Research  
National Marine Fisheries Service, Coral Reef Ecosystem Division  
1125B Ala Moana Blvd., Honolulu, HI 96814

**Abstract**—Seven shallow-water banks, shoals and isolated reefs of the Mariana Archipelago were surveyed for sea turtles using surface and underwater survey techniques as part of efforts to ascertain turtle distributions and numbers throughout the Mariana region. Three green turtles (*Chelonia mydas*) were observed, one each at Supply Reef, Zealandia Bank and Arakane Reef. No other turtle species were seen. Twenty-six species of macroalgae were identified at six of the seven reefs, three (12%) of which are known to be utilized by green turtles. The limited number of turtles at isolated reef systems may be due to low recruitment rates, inadequate habitat range and resources, increased exposure to predation, and/or increased effort required to remain on location. Although sea turtles in some cases utilize shallow isolated reef systems, these systems do not appear to support large turtle numbers on a resident basis in the Mariana region.

### Introduction

Quantitative information on sea turtle abundances and distributions is a prerequisite to understanding the status and recovery of turtle populations within U.S. territorial waters of the Pacific (NMFS & USFWS 1998a-f). The National Marine Fisheries Service has advised, sponsored and/or participated with local agencies in assessing the location and status of resident and breeding turtle populations in Guam and the southern arc islands of the Commonwealth of the Northern Mariana Islands (CNMI; Johannes 1986, McCoy 1997, NMFS & USFWS 1998a-e, Pultz et al. 1999, Belt Collins Hawaii & Marine Research Consultants 2000, Kolinski et al. 2001, 2004, Ilo & Manglona 2001, 2002, Cummings 2001, 2002, Belt Collins Hawaii 2003, The Environmental Company 2004). Although four species of turtles (*Chelonia mydas*, *Eretmochelys imbricata*, *Dermochelys coriacea* and *Lepidochelys olicavea*) have been observed in

Mariana Archipelago waters (Pritchard 1982, NMFS & USFWS 1998a, c), recent estimates suggest between 1,000 and 2,000 mainly immature resident green turtles reside around southern arc islands of the CNMI (Kolinski et al. 2004). Such surveys for resident turtles have recently expanded, with the aid of the NOAA Ship R/V *Oscar Elton Sette*, to encompass the remote northern arc islands along with many isolated reefs throughout the Mariana region. This paper presents information on in-water assessments for turtles at seven isolated reef systems along the northern, southern and back arcs of the Mariana Archipelago. To our knowledge this is the first time these areas have been surveyed for sea turtles.

### Study Area

The Mariana Archipelago consists of northern and southern arc islands and reefs and a back-arc of isolated reefs oriented parallel in a north-south direction from 12.5° to 20.8°N (142.4° to 146.5°E, Figure 1). The area is politically divided between the U.S. Territory of Guam (Guam Island and surrounding reef systems) and the U.S. Commonwealth of the Northern Mariana Islands (comprised of 14 islands extending from Rota to Uracas and regional submerged reef systems). Twenty-eight isolated reef systems (i.e., reefs not associated with land) are reported to occur within the region (Defense Mapping Agency Hydrographic/Topographic Center 1979, Hunter 1995, Micronesian Environmental Services 1997, Western Pacific Regional Fishery Management Council 2002). Half of these appear shallow enough to potentially be suitable for resident sea turtle habitation (< 40 m depth; Defense Mapping Agency Hydrographic/Topographic Center 1979, Micronesian Environmental Services 1997).

Seven isolated reef systems were surveyed: Stingray Shoals (20°29.64'N, 142°26.31'E), Pathfinder Reef (16°30.39'N, 143°8.93'E), and Arakane Reef (15°38.03'N, 142°45.67'E) in the back-arc; Supply Reef (20°8.49'N, 145°5.86'E) and Zealandia Bank (16°53.70'N, 145°51.02'E) in the northern-arc of the CNMI; Tatsumi Reef (14°54.18'N, 145°39.77'E) in the southern-arc of the CNMI; and Santa Rosa Reef (12°49.47'N, 144°25.55'E), located south of Guam (Figure 1). Reef areas and shallowest recorded depths are located in Table 1. Stingray Shoals was dominated by a topographically complex hermatypic scleractinian coral community with steeply sloped shoal walls. Arakane and Pathfinder Reefs were characterized by convoluted mounds of pavement with soft and hard corals. The mounds were separated by wide meandering grooves of sand, rubble and/or pavement. Supply Reef had a 30 m deep plateau with ridges reaching 9 m depth. The reef was rugose and was dominated by hermatypic scleractinian corals. Zealandia Bank had a 33 m deep platform with two thin basalt pinnacles occasionally breaking the waters surface. Platform topography appeared similar to that of Stingray Shoals, with topographic complexity a result of coral accretion. The majority of Tatsumi Reef appeared as pavement with scattered corals. However, breaks within the reef provided limited topographic complexity. Santa Rosa Reef consisted of aggregations of corals and sand, and large areas of pavement.




Figure 1. Map of islands and reported isolated reef systems in the Mariana Archipelago. ★ = isolated reef system. Sites assessed are in bold.

## Methods

All surveys were conducted off the NOAA Ship R/V *Oscar Elton Sette* between 25 August and 28 September 2003. A combination of tow-board, dive and surface surveys were employed in assessing turtle presence. Towed-diver surveys (Hill & Wilkinson 2004) consisted of three SCUBA divers being towed by a single boat. Two of the divers searched from distances roughly 1 to 2 m above the substrate, while a third diver searched for turtles from a shallower depth in the water column. Although tows transited deeper waters, dive depths did not exceed 40 m and covered mainly shallower waters when surveying submerged reef tops and edges. Dive surveys were made between depths of 13 and 25 m by two additional teams of three and five people using SCUBA who collected data on turtles while assessing fish, invertebrate and algal (Preskitt et al. 2004) community composition along three 25 m transects at each dive location (dive surveys at Zealandia Bank were conducted along pinnacle walls where lines could not be laid. Transect lengths were estimated at 50 m for each of two pinnacles). A third team, responsible for mooring unrelated scientific instrumentation, also watched for and reported on turtle sightings at Supply and Santa Rosa Reefs. The greatest lengths of the areas transited underwater by this team were estimated and reported with the dive survey information. Surface waters covering the reefs were transited and searched extensively by three to five support boats at each site. All boat and dive personnel were familiarized with turtle search and reporting objectives and were interviewed by a turtle survey coordinator following each site visit. Turtles observed were identified to species and their approximate locations were marked using Garmin 76S handheld geographic positioning systems. Observation time, turtle activity, size (visually estimated straight carapace length, SCL), and habitat related characteristics were recorded.

Fleshy macroalgae were collected from two survey routes at each isolated reef, with the exception of Supply (one site) and Tatsumi Reefs (no alga collections made), and were identified to species. Coralline algae were lumped into crustose and upright functional groups. Minute turf algae, although present at all sites, were not identifiable in the field and are not considered in this study.

## Results

A total of 34 hrs of submerged and 36 hrs of surface surveys were conducted at the seven isolated reefs systems. This included 21 tows and 13 separate dives, with up to five boats used for surface surveys. A total of only three individual green turtles (*Chelonia mydas*) were observed at the seven reefs: one each at Supply Reef, Zealandia Bank and Arakane Reef. No turtles were observed at Stingray Shoals, Pathfinder Reef, Tatsumi Reef or Santa Rosa Reef. Two of the turtles were of juvenile size (< 70 cm SCL), and one was of a larger size where maturity status was difficult to determine (Table 1).

Table 1. Turtle observations at isolated reef systems of the Mariana Archipelago during 25 August to 28 September 2003 surveys.

Site	Date	Reef Area (km <sup>2</sup> ) within		Shallowest Depth (m)	Method	Time (hr:min)	No. Transects	Transect Length (km)	No. Turtles Observed	Species	Estimated Turtle Size SCL (cm)
		(40 m)	(100 m)								
Stingray Shoals	8/29	0.28	0.67	17	Surface	6:24	(4 boats)	-	0		
					Towed-diver	2:52	4	7.23	0		
					Dive	2:13	2	0.15	0		
Supply Reef	9/1	0.11	0.55	9	Surface	6:59	(5 boats)	-	0		
					Towed-diver Dive	0:27 2:10	1 2	1.16 0.13	0 1	Gn <sup>(b)</sup>	60
Zealandia Bank	8/25	0.09	1.5	0	Surface	5:47	(5 boats)	-	1 <sup>(a)</sup>	Gn	55
					Dive	2:32	2	0.10	1 <sup>(a)</sup>	Gn	55
Pathfinder Reef	9/14	1.01	1.36	12	Surface	6:58	(4 boats)	-	0		
					Towed-diver	3:48	4	9.73	0		
					Dive	2:32	2	0.15	0		
Arakane Reef	9/15	0.72	1.08	10	Surface	8:40	(4 boats)	-	0		
					Towed-diver	5:12	6	13.80	1	Gn	80
					Dive	2:47	2	0.15	0		
Tatsumi Reef	9/17	3.0	6.6	11	Surface	2:32	(5 boats)	-	0		
					Towed-diver	2:43	3	8.72	0		
Santa Rosa Reef	9/28	25.1	30.7	9	Surface	8:26	(3 boats)	-	0		
					Towed-diver	2:49	3	7.38	0		
					Dive	3:44	3	0.18	0		

<sup>(a)</sup> Reef areas within 40 m and 100 m depth contours were determined from boat and shipboard measurements, except Tatsumi and Santa Rosa Reefs, which were based on NOAA charts.

<sup>(b)</sup> The two turtle observations at Zealandia Bank were likely of the same turtle at different times and depths.

<sup>(c)</sup> Gn = green turtles.

Table 2. Algal species located at CNMI and Guam banks, shoals and isolated reefs. Station numbers where alga were found are presented to indicate commonality on a reef.

Alga	Stingray Shoals	Supply Reef	Zealandia Bank	Pathfinder Reef	Arakane Reef	Santa Rosa Reef	Listed Food Source <sup>(1)</sup>
<b>Chlorophyta</b>							
<i>Avrainvillea nigricans</i>						1, 2	
<i>Boodlea composita</i>						2	
<i>Boodlea vanbosseae</i>					1		
<i>Caulerpa peltata</i>				1			
<i>Caulerpa urvilleana</i>						1, 2	✓
<i>Caulerpa</i> sp.						2	? <sup>(2)</sup>
<i>Chlorodesmis</i> sp.				1, 2	1		
<i>Dictyosphaeria cavernosa</i>					2	1, 2	✓
<i>Dictyosphaeria versluysii</i>		1	1, 2	1	1	1, 2	✓
<i>Halimeda cylindracea</i>						1, 2	
<i>Halimeda taenicola</i>			1, 2	1, 2	1, 2	2	
<i>Halimeda</i> spp.			1, 2		1, 2	2	?
<i>Microdictyon okamurai</i>					1, 2	1, 2	
<i>Neomeris</i> sp.				1, 2		1	
<i>Rhipidosiphon javensis</i>					2		
<i>Rhipilia sinuosa</i>			1		2		
<i>Udotea argentea</i>						1, 2	
<i>Ventricaria ventricosa</i>		1	1, 2	1, 2	2		
<b>Phaeophyta</b>							
<i>Lobophora variegata</i>		1					
<b>Rhodophyta</b>							
crustose coralline		1		1, 2	1, 2	1	
gelid						1, 2	?
<i>Laurencia</i> sp.	1, 2			1			?
<i>Liagora ceranoides</i>						1	
non-geniculate branched coralline						1	
<i>Peyssonnelia rubra</i>				2			
<b>Cyanophyta</b>							
Total No.		1	1, 2	1, 2	1, 2	2	
26	1	5	6	10	12	17	3 to 7

<sup>(1)</sup>Species listed as food for green turtles by Hirth (1997).

<sup>(2)</sup>? = Listed as green turtle food by Hirth (1997), but species may differ.

Algal collections from six of the seven reef systems revealed 26 macroalgal species, with an increase in diversity from north to south (Table 2). The number of observed species was positively correlated with reef area ( $\log_{10}$  transformed) within the 40 m contour (Pearsons  $r = 0.845$ ,  $P = 0.034$ ). Green algae accounted for 69% of total species recorded, with species of the calcified genus *Halimeda* being among the most dominant algae at all sites except for Stingray Shoals. When compared to lists of plants known to be eaten by green turtles, only three

(12%) of the species found represented known food sources (Hirth 1997), with limited abundance in areas observed (P. Vroom pers. obs.). However, unidentified species in four recorded genera may also be utilized. Edible algal species were identified on reefs where green turtles were observed (Table 2).

## Discussion

This study is the first we know of to survey for turtles at banks, shoals and other isolated reefs within the Mariana Archipelago and is a part of ongoing efforts to ascertain the abundance and distribution of resident turtles throughout the CNMI (Pultz et al. 1999, Belt Collins Hawaii & Marine Research Consultants 2000, Kolinski et al. 2001, 2004, in prep., Ilo & Manglona 2001, 2002, Belt Collins Hawaii 2003, The Environmental Company 2004) and the Territory of Guam (Davis et al. unpubl. data, Kolinski et al. unpubl. data). The seven sites are likely to be representative of shallow (< 40 m) isolated reef systems throughout the archipelago. The absence or limited number of turtle sightings is consistent with reports by fishers visiting these and other isolated reefs in the region (Larry Ilo, CNMI Division of Fish & Wildlife, pers. comm., Gerry Davis, Guam Division of Aquatic & Wildlife Resources, pers. comm.).

Surveys to date have highlighted the numerical dominance of green turtles within Mariana waters (Pritchard 1982, Wiles et al. 1995, Pultz et al. 1999, Belt Collins Hawaii & Marine Research Consultants 2000, Ilo & Manglona 2001, 2002, Kolinski et al. 2001, 2004, in prep., Belt Collins Hawaii 2003, The Environmental Company 2004). Green turtles were the only species encountered in this study. The limited numbers or absence of green turtles at these isolated reef systems may be due to a number of factors including: a long history of hunting and egg gathering at Indo-West Pacific breeding rookeries; low probability of turtles transitioning from the post-hatchling pelagic phase encountering small isolated reefs for benthic recruitment; inadequate habitat and forage resources; increased exposure to predation by sharks, and; increased energetic effort required to remain over a small isolated reef during high ocean wave and current events. Observations of low diversity and abundance of listed (Hirth 1997) green turtle food resources within the limited areas surveyed suggest that, while green turtles appear to be supported in low numbers at certain isolated reefs, perhaps for transitory periods of time, overall resources may be inadequate to support large resident populations.

Although the full contribution of isolated reef systems to turtle population status and dynamics remains unknown, these systems do not presently appear to support large turtle numbers on a resident basis in the Mariana region. This information has obvious implications when attempting to project turtle population sizes for the CNMI (Kolinski et al. 2004, in prep.), and Territory of Guam (Davis et al. unpubl. data, Kolinski et al. unpubl. data). Unless a seasonal component exists, these low numbers suggest that interactions between sea turtles and bottom-fishing activities in these areas are likely to be rare. Research, management

and recovery efforts might best be focused around island areas where turtles are known to occur in larger numbers and impacts are likely to be identified and subsequently addressed.

### Acknowledgements

Special thanks to Captain Ken Barton and the crew of the NOAA Ship *Oscar Elton Sette* for the voyage and survey support. Personnel that participated in collecting data on turtles included Joe Laughlin, Brian Zgliczynski, Molly Timmers, Rusty Brainard, Robert Schroeder, Mike Trianni, Kate Moots, Fran Castro, Peter Houk, Machel Malay, Trina Leberer, Shawn Wusstig, Brent Tibbatts, Mark Capone, and Doug Roberts. Richard Seman, Ray Clarke, and Charles Karnella were instrumental in prioritizing turtle surveys as a cruise objective. John Naughton and Alan Everson provided program support. Local support and information provided by Larry Ilo and the CNMI Division of Fish & Wildlife, and Gerry Davis and the Guam Division of Aquatic and Wildlife Resources is greatly appreciated. Thanks to George Balazs, Ray Clarke, Charles Karnella, Evelyn Cox, Samuel Pooley and an anonymous reviewer for suggestions which improved the manuscript.

### References

- Belt Collins Hawaii. 2003. Year 2002 assessment of marine and fisheries resources, Farallon de Medinilla, Commonwealth of the Northern Mariana Islands. Final report prepared for the Department of the Navy, Pacific Division, Naval Engineering Command, Pearl Harbor, Hawaii, Contract No. N62742-99-D-1807. 62 pp.
- Belt Collins Hawaii & Marine Research Consultants. 2000. Year 2000 assessment of marine and fisheries resources, Farallon de Medinilla, Commonwealth of the Northern Mariana Islands. Final report prepared for the Department of the Navy, Pacific Division, Naval Engineering Command, Pearl Harbor, Hawaii. 45 pp.
- Cummings, V. 2001. Guam sea turtle recovery report two. Division of Aquatic & Wildlife Resources Report, Guam Department of Agriculture, Mangilao, Guam. 5 pp.
- Cummings, V. 2002. Guam sea turtle recovery report three. Division of Aquatic & Wildlife Resources Draft Report, Guam Department of Agriculture, Mangilao, Guam. 9 pp.
- Defense Mapping Agency Hydrographic/Topographic Center. 1979. Loran C north Pacific Ocean Mariana Islands—Guam, Marianas surface/air operating areas. Map 81025. Washington D.C.
- Hill, J. & C. Wilkinson. 2004. Methods for ecological monitoring of coral reefs, a resource for managers. Version 1. Australian Institute of Marine Science, Townsville, MC, QLD, Australia. 117 pp.



- Hirth, H. F. 1997. Synopsis of the biological data on the green turtle *Chelonia mydas* (Linnaeus 1758). Fish and Wildlife Service Biological Report 97(1). 120 pp.
- Hunter, C. L. 1995. Review of status of coral reefs around American Flag Pacific Islands and assessment of need, value and feasibility of establishing a coral reef fishery management plan for the western Pacific region. Final report prepared for the Western Pacific Regional Fishery Management Council, Honolulu, Hawaii. 30 pp.
- Ilo, L. I. & J. M. Manglona. 2001. Rota turtle assessment report. Report prepared for the Division of Fish and Wildlife, Saipan, Commonwealth of the Northern Mariana Islands. 22 pp.
- Ilo, L. I. & J. M. Manglona. 2002. Sea turtle assessment report for the island of Anatahan. Report prepared for the Division of Fish and Wildlife, Saipan, Commonwealth of the Northern Mariana Islands. 16 pp.
- Johannes, R. E. 1986. A review of information on the subsistence use of green and hawksbill sea turtles on islands under United States jurisdiction in the western Pacific ocean. NMFS Southwest Region Administrative Report SWR-86-2. 41 pp.
- Kolinski, S. P., L. I. Ilo & J. M. Manglona. 2004. Green turtles and their marine habitats at Tinian and Aguijan, with projections on resident turtle demographics in the southern arc of the Commonwealth of the Northern Mariana Islands. *Micronesica* 37: 95–116.
- Kolinski, S. P., D. M. Parker, L. I. Ilo & J. K. Ruak. 2001. An assessment of the sea turtles and their marine and terrestrial habitats at Saipan, Commonwealth of the Northern Mariana Islands. *Micronesica* 34: 55–72.
- McCoy, M. A. 1997. The traditional and ceremonial use of the green turtle (*Chelonia mydas*) in the Northern Mariana Islands with recommendations for its use in cultural events and education. Final report for the Western Pacific Regional Fishery Management Council and the University of Hawaii Sea Grant College Program, Honolulu, Hawaii. 87 pp.
- Micronesian Environmental Services. 1997. The Commonwealth of the Northern Mariana Islands: an assessment of the coral reef resources under local and federal jurisdiction. Final report prepared for the Western Pacific Regional Fishery Management Council, Honolulu, Hawaii, Contract No. 96-WPC-119. 69 pp.
- NMFS & USFWS. 1998a. Recovery plan for U.S. Pacific populations of the green turtle (*Chelonia mydas*). National Marine Fisheries Service, Silver Spring, MD. 84 pp.
- NMFS & USFWS. 1998b. Recovery plan for U.S. Pacific populations of the hawksbill turtle (*Eretmochelys imbricata*). National Marine Fisheries Service, Silver Spring, MD. 82 pp.
- NMFS & USFWS. 1998c. Recovery plan for U.S. Pacific populations of the olive ridley turtle (*Lepidochelys olivacea*). National Marine Fisheries Service, Silver Spring, MD. 52 pp.

- NMFS & USFWS. 1998d. Recovery plan for U.S. Pacific populations of the leatherback turtle (*Dermochelys coriacea*). National Marine Fisheries Service, Silver Spring, MD. 65 pp.
- NMFS & USFWS. 1998e. Recovery plan for U.S. Pacific populations of the loggerhead turtle (*Caretta caretta*). National Marine Fisheries Service, Silver Spring, MD. 59 pp.
- NMFS & USFWS. 1998f. Recovery plan for U.S. Pacific populations of the east Pacific green turtle (*Chelonia mydas*). National Marine Fisheries Service, Silver Spring, MD. 50 pp.
- Preskitt, L. B., P. S. Vroom & C. M. Smith. 2004. A rapid ecological assessment (REA) quantitative survey method for benthic algae using photoquadrats with SCUBA. *Pacific Science* 58: 201–209.
- Prichard, P. C. H. 1982. Marine turtles of Micronesia. In K. A. Bjorndal (ed), *Biology and Conservation of Sea Turtles*, pp. 263–274. Smithsonian Institution Press, Washington D.C.
- Pultz, S., D. O'Daniel, S. Krueger, H. McSharry & G. Balazs. 1999. Marine turtle survey on Tinian, Mariana Islands. *Micronesica* 32: 85–94.
- The Environmental Company. 2004. Year 2003 assessment of marine and fisheries resources, Farallon de Medinilla, Commonwealth of the Mariana Islands. Final report prepared for the Pacific Division, Naval Facilities Engineering Command, Pearl Harbor, Hawaii, Contract No. N62742-02-D-1802. 46 pp.
- Western Pacific Regional Fishery Management Council. 2002. Options for managing the bottomfish fishery in federal waters around Guam, for consideration in adjusting or amending the fishery management plan for the bottomfish and seamount groundfish fisheries of the western Pacific region. Draft report, Western Pacific Regional Fishery Management Council, Honolulu, Hawaii. 37 pp.
- Wiles, G. J., C. F. Aguon, G. W. Davis & D. J. Grout. 1995. The status and distribution of endangered animals and plants in northern Guam. *Micronesica* 28: 31–49.