

Ramirez-Cruz, J.C., Ramirez, I.P. and Flores, D.V. 1991. Distribución y abundancia de la tortuga perica, *Caretta caretta* en la costa occidental de Baja California Sur, Mexico. Archelon 1:1-4.

Villanueva, D. 1991. La tortuga perica *Caretta caretta gigas* (Deraniyagala 1939) en la costa de Pacifico de Baja California Sur, Mexico. Departamento de Biología Marina. Univ. Autónoma de Baja California Sur, La Paz, BCS, Mexico. Tesis de Licenciatura. 68 p.

## Movement and dive behavior determined by satellite telemetry for male and female olive ridley turtles in the Eastern Tropical Pacific

Denise M. Parker<sup>1</sup>, Peter H. Dutton<sup>2</sup>, Kerry Kopitsky<sup>2</sup>, and Robert L. Pitman<sup>1</sup>

<sup>1</sup> Joint Institute for Marine and Atmospheric Research, 8607 La Jolla Shores Drive, P.O. Box 241, La Jolla, California 92037, USA

<sup>2</sup> NOAA, National Marine Fisheries Service, Southwest Fisheries Science Center, 8607 La Jolla Shores Drive, P.O. Box 241, La Jolla, California 92037, USA

<sup>3</sup> College of Marine Studies, University of Delaware, Robinson Hall, Newark, Delaware 19717, USA

Olive ridleys have a worldwide distribution in mainly tropical areas. Nesting occurs along continental margins in all oceans. The largest nesting areas in the Eastern Pacific are along the coasts of Mexico and Costa Rica. Many studies have been done on nesting populations and movement from the nesting grounds, but few studies have been done on the movement of olive ridleys in the open ocean (Plotkin et al. 1994, Beavers and Cassano 1996). This paper summarizes the movement and dive behavior of male and female olive ridleys tracked during 1999-2000.

Satellite-linked Time-Depth Recorders (SDRs, Wildlife Computers) were deployed on eight olive ridley turtles, four adult females, three adult males, and one juvenile, during the 1999 *Stenella* Abundance Research Survey (STAR) in the Eastern Tropical Pacific. Turtles were spotted from NOAA ships traveling set course lines. Small boats were deployed to capture turtles rodeo style by hand, and turtles were brought back to the ship for measurements and satellite attachment. The SDRs were attached using fiberglass cloth and polyester resin based on the methods by Balazs et al. (1996). Dive data from the SDRs were compiled using Wildlife Computers SatPak30 and duration of dives and time-at depth were examined. The time-at-depth bins recorded the amount of time spent at each preset depth. Three of initial SDRs were programmed with parameters starting at 1-meter (m) depth (depth histogram bins: 1, 5, 10, 15, 20, 25, 30, 40, 50, 60, 80, 100, 150, 150+ meters). However, since an objective for this on-going study was to be able to link the dive data with visual surveys, the parameters were changed. The final five SDRs were programmed to include a bin that specifically counted time on surface (depth histogram bins: 0, 1, 5, 10, 15, 20, 30, 40, 50, 60, 80, 100, 150, 150+ meters). However, for purposes of analysis in this presentation, the 0 and 1 meter depth bins were combined for these five SDRs so the data could be compared to the previous three. T-test analyses were done to determine significant differences in dive behaviors between male and female turtles.

The movements of all of the olive ridleys are seen in Fig. 1. The three mated female olive ridleys moved directly to or offshore of a major nesting area. The two males captured mating with two of the females were seen to continue to move along the coast. The non-mating female and male both stayed offshore. While all of the turtles recorded a high percentage of short dives, 60% of the dives were 2 min or less in duration, table 1 shows the maximum dive duration for each turtle. The average of the longest dive time for females was 120-180 minutes, 75 minutes for males, and 45-60 minutes for the one juvenile. There were no significant difference between male and females for any of the dive duration bins.

Visual observations of olive ridleys at sea have indicated that they may spend a large amount of their time on the surface. Average time-at-surface calculated for the five SDRs that recorded 0-m data indicated surface times of 10-22%, and even with depths to 1-m included, times near the surface were still low, 19-44%. However, daily, diurnal and variations between

turtles were seen. Daily time-near surface averages ranged between 10 and 50%, and this varied between turtles. A diurnal dive behavior was seen where most turtles spent more time near the surface during daylight hours, which were between 9am-2 pm, between 22-56% (mean of 37%) of the total dive time was spent near the surface during this 6-hour period. There was no difference seen for time spent near the surface between male and female turtles.

There were no significant differences in time-at-depth and dive duration between turtles that had mated and not mated. Although all turtles had the opportunity to make deep dives, the mated females and males did not make dives greater than 150m, even while over open ocean. The non-mated, pelagic male and female both made dives greater than 150m with a number of dives over 200m, some which registered at the limit that the SDR could record (250m). On the other hand, a mated female that had moved inshore and possibly nested spent 6 months off the coast of Nicaragua. During that time, it stayed in waters that were less than 150m in depth and made regular dives to depths that would have reached the bottom - possibly either resting or foraging on the bottom.

Fig. 2 shows the average time-at-depth profiles for female and male olive ridley turtles. Females spent significantly more time at 40 and 80m depths than did males. It has been shown by Donguy and Meyers (1987) and also Fiedler (1992), that within the Eastern Tropical Pacific there is a permanent thermocline between 20 and 100m (around the 20° C isotherm). All of our adult turtles spent significant amounts of time, at least 25% of total dive time, in the area of the thermocline.

An SDR transmitter put on only one juvenile turtle unfortunately had a very short track, only 5 days in length. The juvenile stayed near shore in waters that were between 100 - 200m in depth. During the track, the daily and diurnal differences in time spent near surface were similar to that of the adult turtles, but there were visual differences between the juvenile and adult dive profiles. The juvenile made no dives greater than 80m even though it was in water where deeper dives could be made, and 75% of its time was spent within 10m of the surface.

In conclusion, the low average times that the turtles spent near the surface were a combination of diurnal and daily differences in dive behavior. Female olive ridleys in this study spent significantly more time at 40 and 80 meters than did the males, and the thermocline is an important foraging area for the olive ridley as both male and female turtles spent a significant amount of time in the region of the thermocline - between 20-100m. The significance of these findings are that the data obtained for time at surface when taking into account the diurnal variability will be able to be combined with observers visual counts to gain a better estimate of pelagic numbers of olive ridleys in the Eastern Tropical Pacific. Also the information on dive depths and behavior can be also be used as a tool to help mitigate and reduce bycatch in fisheries.

LITERATURE CITED

Balazs, G.H., R.K. Miya, and S.C. Beavers. 1996. Procedures to attach a satellite transmitter to the carapace of an adult green turtle, *Chelonia mydas*. In: J.A. Keinath, D. Bernard, J.A. Musick, and B.A. Bell, comps. Proc. of the Fifteenth Annual Symposium on Sea Turtle Biology and Conservation, NOAA Tech. Memo. NMFS-SEFSC-387. p. 21-26.

Beavers, S.C. and E.R. Cassano. 1996. Movements and dive behavior of a male sea turtle (*Lepidochelys olivacea*) in the eastern tropical Pacific. *Journal of Herpetology* 30:97-104.

Donguy, J.R. and G. Meyers. 1987. Observed and modeled topography of the 20°C isotherm in the tropical Pacific. *Oceanol. Acta* 10:41-48.

Fiedler, P.C. 1992. Seasonal climatologies and variability of eastern tropical Pacific surface waters. NOAA Tech. Rep., NMFS 109.

Plotkin, P.T., R.A. Byles, and D.W. Owens. 1994. Post-breeding movements of male olive ridley sea turtles *Lepidochelys olivacea* from a nearshore breeding area. In: K.A. Bjorndal, A.B. Bolten, D.A. Johnson, P.J. Ellazar, comps. Proceedings of the Fourteenth Annual Symposium on Sea Turtle Biology and Conservation, NOAA Tech. Memo. NMFS-SEFSC-351. p. 19.

Table 1. Maximum dive duration recorded for four female, three male, and one juvenile olive ridley turtles. Data were received from satellite time-depth recorders attached to each turtle before release during the 1999 Stenella Research Cruise in the Eastern Tropical Pacific.

Turtle ID	Sex	Maximum Dive Duration
21128	Female	60 - 75 minutes
21130	Female	120 - 180 minutes
21136	Female	180 - 240 minutes *
24645	Female	120 - 180 minutes
21129	Male	75 - 90 minutes *
21138	Male	60 - 75 minutes
21143	Male	60 - 75 minutes
24644	Juvenile	45 - 60 minutes




Fig. 1. Map of the movement of eight olive ridley turtles (four female, three male, and one juvenile) that were released with satellite time-depth recorders during the 1999 Stenella Research Cruise in the Eastern Tropical Pacific.




Fig. 2. Average time-at-depth dive profiles for four female (left) and three male (right) olive ridleys during 1999-2000. Time-at-depth was averaged for each depth bin over the complete track of an individual turtle (one bar indicates one turtle). Significant differences ( $p < 0.05$ ) were found between male and female turtle's time-at-depth for the 40 and 80-m depth bins shown by an asterisk.



NOAA Technical Memorandum NMFS-SEFSC-503

PROCEEDINGS  
OF THE TWENTY-SECOND ANNUAL  
SYMPOSIUM ON SEA TURTLE  
BIOLOGY AND CONSERVATION



**M i a m i . U S A**

4 to 7 April 2002  
Miami, Florida, USA

Compiled by:  
Jeffrey A. Seminoff

U.S. Department of Commerce  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southeast Fisheries Science Center  
75 Virginia Beach Drive  
Miami, FL 33149 USA

August 2003



NOAA Technical Memorandum NMFS-SEFSC-503

PROCEEDINGS  
OF THE TWENTY-SECOND ANNUAL  
SYMPOSIUM ON SEA TURTLE  
BIOLOGY AND CONSERVATION

4 to 7 April 2002  
Miami, Florida, USA

Compiled by:

Jeffrey A. Seminoff

U.S. DEPARTMENT OF COMMERCE  
Donald L. Evans, Secretary

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
Conrad C. Lautenbacker, Jr., Administrator

NATIONAL MARINE FISHERIES SERVICE  
William T. Hogarth, Assistant Administrator for Fisheries

Technical Memoranda are used for documentation and timely communication of preliminary results, interim reports, or special-purpose information, and have not received complete formal review, editorial control or detailed editing.