



ISSN: 2993-8864

DOI: 10.33552/OJEEES.2025.02.000539

Online Journal of
Ecology & Environment Sciences

Iris Publishers

Review Article

Copyright © All rights are reserved by Roxane Tran

Prevalence of Fibropapillomatosis in Sea Turtles of New Caledonia

Roxane Tran^{1*}, Claire Goiran², Jean-Jérôme Cassan³, Frédéric Avril³, Izabella Lach⁴, Agathe Binois⁵, Laurent Fabre⁶, Olivier Chateau⁷ and Tyffen Read⁸.

¹National Institute of Applied Sciences, 31400, Toulouse, France, **ORCID:** 0009-0001-1358-8697

²University of New Caledonia, Nouméa, New Caledonia. **ORCID:** 0000-0002-1649-4418

³Direction du développement économique et de l'environnement, North Province, 98860, Koné, New Caledonia. **ORCID:** 0000-0002-2757-0623

⁴Clinic "Les 3 cliniques", 98800, Nouméa, New Caledonia

⁵Clinic "Les palmiers", 98819, Moindou, New Caledonia

⁶Clinic "Clinique Vétérinaire du Récif", 98850, Koumac, New Caledonia

⁷Laboratory of marine biology and ecology, Aquarium des Lagons, 98800, Nouméa, New Caledonia. **ORCID:** 0000-0003-1153-6284

⁸Direction du Développement Durable des Territoires, South Province, 98800, Nouméa, New Caledonia. **ORCID:** 0000-0002-4385-0043

Corresponding author: Roxane Tran, National Institute of Applied Sciences, 31400, Toulouse, France

Received Date: August 07, 2025

Published Date: October 08, 2025

Abstract

This study investigated the presence of tumors in sea turtles of New Caledonia, which were confirmed to be fibropapilloma (FP) lesions by necropsies. The data presented here was gathered from several projects, from 2011 to 2025, using different collecting methods such as stranding reports (44%), photo-identification (40%), poaching (8%), Capture-Mark-Recapture study (4%) and traditional hunting (4%). Among the three species of sea turtles regularly seen in New Caledonian waters, *Chelonia mydas* represented 100% of the turtles affected by the disease. The great majority of the reports were recorded in South province (72%), the most populated area of New Caledonia. Nearly half of the affected individuals were stranded juveniles. Photo-identification allowed the detection of the disease on almost half of the affected individuals and also the monitoring of the evolution of the tumors which were seen growing in most of the cases. This method confirmed regression of FP for one turtle. FP cases appear to be slowly increasing in New Caledonia, so continued monitoring to see if this trend persists is advisable.

Key Words: Green turtles; Fibropapilloma; Photo-identification; South Pacific; Stranding



Introduction

Fibropapilloma is a tumor disease that affects all seven species of sea turtles, with a higher prevalence in green turtles (*Chelonia mydas*) (Whilde et al. [1]). It was first reported in green turtles from the state of Florida, southern United States in 1938 (Smith and Coates [2]). Fibropapilloma tumors are found on the skin and occasionally on the upper and lower shell of sea turtles. External tumors are histologically benign but can be fatal depending on their size and distribution (Page-Karjian [3]). When tumors are small, there is a minimum threat of life to the affected turtle. However, bigger tumors can disrupt locomotion, foraging ability, vision, and predator evasion (Herbst [4]). In the case of internal tumors, they affect vital functions and lower chances of survival of sea turtles (Whilde et al. [1]).

A herpes virus, chelonian herpes virus (ChHV5), is closely associated with the presence of tumors (Jones et al. [5]). Even though the virus has coexisted with turtles for millions of years (Whitlock et al. [6]), the frequent reporting of the disease in turtles that reside in poor-quality eutrophicated waters suggests that the disease could have an environmental component (Herbst and Klein [7]). Possible routes of transmission of FP include leeches (*Ozobranchus* spp.), cleaner fishes and direct contact (Lu et al. [8]; Greenblatt et al. [9]; Work et al. [10]).

A growing number of scientific articles have been published over the last two decades reporting fibropapilloma cases in new geographical areas or describing current local coastal prevalence (Dupont et al. [11]). Fibropapilloma was first seen in the Atlantic (Smith and Coates [2]), but since has been documented in multiple areas in the Pacific (Greenblatt et al. [12]), Indian Ocean (Jones et al. [22]), Asia (Adnyana et al. [13]; Lucero et al. [14]; Li et al. [15]; Robben et al. [16]) and Australia (Flint et al. [17]; Ariel et al. [18]; Jones et al. [19]).

New Caledonia is a French archipelago located in the South Pacific. It is divided into three provinces (South, North and Loyalty Islands). The main island is composed of the two first provinces. The South, North and Loyalty Islands Provinces are respectively inhabited by 75, 18 and 7% of the population (ISEE 2019). The main island displays one of the largest barrier-reefs in the world, isolating a locally-wide lagoon (Coudray [20]). Salinity is generally close to ocean salinity but in protected bays it may decrease due to occasional rainfalls or increase during periods of droughts which are especially pronounced during El Niño – Southern Oscillation (ENSO) events (Ouillon et al. [21]). ENSO phenomena can also increase the disequilibrium of oxygen's abundance because temperatures can rise up to 30°C (Rougerie [22]). New Caledonia is geologically underlain by a metalliferous oceanic basement. About one-third of the land surface of the main island consists of massifs containing peridotite (Xun [23]). New Caledonia is currently the third largest producer of nickel in the world (USGS 2023). However, the open cast mining on the hilltops has resulted in major alterations to landscape and habitat strongly enhancing natural soil erosion and subsequent river transport of terrigenous material and associated metals toward the lagoon (Bird et al. [24]). Although all pelagic and benthic communities are affected to some extent by the increases in sediment deposition and decreases in light penetration, fringing reefs are especially sensitive to these

changes (Ouillon et al. [25]).

In New Caledonian waters, three species of sea turtles are seen foraging on a regular basis: green turtles *Chelonia mydas*, loggerhead turtles *Caretta caretta* and hawksbill turtles *Eretmochelys imbricata*; but only the first two are known to nest there (Read et al. [26]). Reefs of New Caledonia host a large number of nesting colonies. Among them, D'Entrecasteaux, Bellona and Chesterfield Reefs shelter the largest green turtle nesting population of the South Pacific (Fretey et al. [27]). New Caledonia also presents well-known loggerhead turtles nesting sites. The most important ones are located on the west coast of the mainland, on La Roche Percée beach, which represents the second highest density nesting site in the South Pacific (Barbier et al. [28]) when other smaller nesting sites were inventoried on dispersed coral islets along the coastline from far north to the 'Grand Lagon Sud' (GLS) at the meridional end of the mainland (Bourgogne et al. [29]).

Chelonia mydas and *Caretta caretta* are known to migrate across the Coral Sea between New Caledonia and Australia (Limpus et al. [30]) where fibropapilloma is enzootic since 2008 (Jones et al. [21]). Identification of large-scale movements was also enabled for individuals recaptured at separated nesting beaches or foraging grounds proving the migratory paths through the Coral Sea (Read et al. [31]). Green turtles are known to migrate back to their foraging site after nesting and thus can cross international waters (Cheng [32]). In New Caledonia, important foraging sites were found in the GLS area (Read et al. [33]), one of six areas of New Caledonia added to the World Heritage List in 2008 (UNESCO).

Fibropapilloma was first documented in New Caledonia in 2011 but no analyses were done to confirm presence of ChHV5, a needed requirement to confirm this disease in a new location (Work and Balazs [34]). The first confirmation of the presence of the chelonid herpesvirus was established in 2014 but was limited to one individual (Read et al. [27]). This study expands on these initial findings and documents additional cases of fibropapilloma in New Caledonia sea turtles using both histology and molecular methods as well as reports of individuals presenting fibropapilloma lesions.

Materials and Methods

Data in this study originated from multiple projects and collecting methods. All individuals which presented pathologies or abnormalities were recorded (including suspicions of fibropapilloma). Biological data (species, sex, presence of fibropapilloma) and curved carapace length (CCL) were recorded when possible. For all methods, turtles were pooled into five categories depending on their CCL used as a proxy for approximate age: post-hatchlings (>5...<-35 cm), juveniles (>35...<-65cm), sub-adults (>65...<-90cm), adults (>90 cm) and unknown (Limpus et al. [35]).

Fieldwork conducted between 2012 and 2014 in the GLS area, allowed the detection of affected turtles using the Capture-Mark-Recapture (CMR) method. These sampling campaigns used the turtle rodeo method (Limpus et al. [36]) to capture sea turtles (N=183) before marking them with a titanium tag and releasing them into the ocean (Read et al. [33]). Starting in January 1999, stranding of sea turtles in New Caledonia (N=406) were recorded

by the Aquarium des Lagons and the different provinces (Read et al. [37]). All turtles found alive were released and tagged after receiving surgery for the removal of found tumors by qualified veterinarians. Since 2019 (and still going), the growing interest of amateurs in wildlife-research has been valuable for scientists. The “Fantastic Grandmothers” (citizen scientist group) initiated a photo-identification (PID) program in the area of Nouméa (Goiran and Shine [38]) on multiple species including sea turtles (N=1533). Photographs were added to the website inaturalist.

Turtles were assigned a subjective tumor score ranging from 0 (no visible external tumors) to 3 (heavily tumored) that indicated the severity of fibropapilloma (Work and Balazs [39]). This approach relies on a subjective index established by Balazs [40] to assess the degree of severity of external fibropapilloma lesions in sea turtles (Balazs [40]).

For a subset of turtles, tumors were collected in formalin and

90% ethanol. For histology, tumors were processed as described and examined for morphologic characteristics of fibropapillomatosis (Herbst et al. [41]). To confirm presence of ChH5, DNA was extracted from ethanol-preserved tumors using Quiagen DNAeasy kits and PCR for the POL gene done as described

Results

Twenty-five green turtles were reported with fibropapilloma lesions, four were captured, ten were identified by photographs and eleven stranded, from 2011 to 2025 with seven cases in North province and the remainder in South province (Figure 1). The ten turtles with FP identified by photos were from the South province, in Nouméa or Bourail (Table 1). Of turtles that could be aged, nine were juveniles, two were sub-adult, three were adults and no age class information was available for one specimen. Of twenty-three turtles assigned a tumor score, 61% were score 1, 35% score 2 and 4% score 3 (Table 1).



Figure 1: Map of recorded sea turtles affected by fibropapilloma in New Caledonia.

Table 1: Chart of turtles with fibropapilloma reported in New Caledonia.

| Date | Lesions/ Tumor score () | Place of finding | Species | Age class | CCL (cm) | Primary activity | Collecting method | Additional comments |
|----------------|-------------------------------|------------------------------|-----------------------|-----------|----------|------------------|---------------------|---|
| June/2011 | Lower shell (2) | Poum, Tiabet/ North Province | <i>Chelonia mydas</i> | Unknown | N/A | Stranded | Stranding | |
| June/2013 | Both eyes and fins (2) | Koumac/ North Province | <i>Chelonia mydas</i> | Sub-adult | 79.5 | Stranded (dead) | Stranding | |
| January/2014 | Fin (1) | Ouegoa/ North Province | <i>Chelonia mydas</i> | Adult | N/A | Feeding | Traditional hunting | |
| July/2014 | Both eyes (1) | Ile Ouen/ South Province | <i>Chelonia mydas</i> | Juvenile | 53.5 | Feeding | CMR | Released after surgery - PCR and histology done: confirmed positive |
| September/2016 | Both eyes and lower shell (2) | Koumac/ North Province | <i>Chelonia mydas</i> | Adult | 92.5 | Stranded (dead) | Stranding | |

| | | | | | | | | |
|----------------|--|--|-----------------------|-----------|------|----------------------------|-----------|--|
| December/2016 | Both eyes, fins, neck, lower shell (2) | Poum/ North Province | <i>Chelonia mydas</i> | Juvenile | 44.5 | Stranded (alive) | Stranding | Released after surgery |
| August/2018 | Both eyes (1) | Ile Ouen/ South Province | <i>Chelonia mydas</i> | Juvenile | N/A | Stranded (dead) | Stranding | PCR done: confirmed positive |
| March/2019 | Both eyes, lower shell, fins (2) | La Foa/ South Province | <i>Chelonia mydas</i> | Juvenile | 60 | Stranded (dead) | Stranding | PCR done: confirmed positive |
| March/2019 | N/A | Poya/ South Province | <i>Chelonia mydas</i> | Sub-adult | 72.5 | Feeding | Poaching | |
| October/2021 | Both eyes, fins (2) | Paita/ South Province | <i>Chelonia mydas</i> | Adult | 93 | Stranded (dead) | Stranding | PCR done: confirmed positive |
| December/2021 | Left eye (1) | Nouméa, Ilot Signal/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID July 2019: "IS24" |
| January/2022 | N/A | Ouegoa/ North Province | <i>Chelonia mydas</i> | Juvenile | 46 | Stranded | Stranding | |
| August/2022 | Left eye (2) | Bourail/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | |
| September/2022 | Both eyes, lower shell, fins (3) | Poya/ South Province | <i>Chelonia mydas</i> | Juvenile | 50 | Feeding | Poaching | Released after surgery |
| November/2022 | Both eyes (1) | Bourail/ South Province | <i>Chelonia mydas</i> | Juvenile | 49 | Stranded (alive) | Stranding | Died before surgery |
| January/2024 | Left eye (1) | Nouméa Ilot Maitre/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID January 2024, "Im650" |
| March/2024 | Right eye (1) | Nouméa, Baie des Citrons/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID December 2023: "Hope". https://www.inaturalist.org/observations/202501597 |
| May/2024 | Right eye (1) | Nouméa, Baie des Citrons/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID September 2020: "Yana". https://www.inaturalist.org/observations/213091478 |
| July/2024 | Left eye (1) | Nouméa/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID July 2024: @684ada9e9fbb. https://www.inaturalist.org/observations/232208015 |
| July/2024 | Left eye (1) | Nouméa/ South Province | <i>Chelonia mydas</i> | Adult | 89 | Stranded (dead) | Stranding | |
| September/2024 | Right eye (1) | Nouméa, Anse Vata/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID September 2024: "Tatianna" |
| November/2024 | Both eyes (1) | Nouméa, Anse Vata/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID January 2021: "Royale" |
| November/2024 | Both eyes and fin (1) | Nouméa/ South Province | <i>Chelonia mydas</i> | Juvenile | 51.3 | Stranded (alive) | Stranding | Released after surgery: "Danielle", first PID November 2022 - PCR and histology done: confirmed positive |

| | | | | | | | | |
|-----------|--------------------|---|---------------------------|----------|-----|---------------------------|-----------|--|
| May/2025 | Left eye (1) | Nouméa, Tepava/ South Province | <i>Chelonia mydas</i> | Unknown | N/A | Feeding | PID | First PID June 2022: "Roxane". https://www.inaturalist.org/observations/283557047 |
| June/2025 | Lower shell (2) | Poum/ North Province | <i>Chelonia mydas</i> | Juvenile | 45 | Stranded (dead) | Stranding | |

Discussion

The first report of an individual carrying suspected lesions of fibropapilloma was recorded in 2011 on a stranded green turtle in the North province, near Tiabet (extreme North of New Caledonia) (Table 1). In 2013, a publication mentioned an individual stranded with fibropapilloma in New Caledonia, but no analysis was done at the time to confirm the virus (Work and Balazs [34]). In 2014, a specimen was found with suspected lesions in South province and a sample was sent to the National Wildlife Health Center-Honolulu field station in Hawaii. Gross and microscopic pathology results were consistent with fibropapillomatosis. Molecular analyses confirmed the presence of chelonid herpesvirus 5 DNA for herpesvirus POL gene on this individual (Read et al. [27]). Five other samples from more recent individuals (Table 1) were analyzed in 2024 and confirmed that all individuals assessed were positive for herpesvirus 5.

A higher number of individuals were reported in South province (72%) than in North province (28%), all methods combined and no individuals were reported in the Loyalty province. This may be due to the higher human population density in South province likely leading to increased chances of reporting of diseased turtles. Indeed, the CMR and photo-identification program were only located in the South province. A large number of stranded animals in South province could be explained by an increasing number of boat owners as well as an escalation of maritime traffic in the last decade who would also be more likely to report stranded turtles.

All turtles found affected by the fibropapilloma virus in this study were green turtles. This concurs with multiple studies around the world that this disease mainly affects green turtles (Smith and Coates [41]; Adnyana et al. [13]; Greenblatt et al. [12]; Li et al. [15]; Jones et al. [19,22]; Robben et al. [16]). In addition, green turtles are the most common species of sea turtle in foraging grounds of New Caledonia which could explain the results of this study (Read et al. [26]; Read and Jean [42]).

The majority of the specimens affected were in growth stages between juveniles and sub-adults. This result could be explained by the importance of these life-stages in the New Caledonian foraging grounds (Read et al. [33]) and concurs with multiple studies worldwide that show that as they enter neritic foraging areas after completing their pelagic life stage, juvenile turtles are at a higher risk of developing fibropapilloma (Ene et al. [43]; Jones et al. [5]; Page-Karjian [3]).

The number of individuals recorded with fibropapilloma

lesions has appeared to increase in recent years with 56% of turtles affected since 2022 (Table 2). How significant this is remains unclear as there could be sampling biases. For instance, the CMR was only done between 2012 and 2014 and was not resumed since and the photo-identification program was initiated in 2021. A more systematic and consistent monitoring effort for stranded and in-water studies might help clarify whether FP is truly increasing or not in New Caledonia.

Data shows that the majority of the turtles with fibropapillomatosis come from stranding reports (44%) (Table 1), establishing the fact that a solid stranding network is needed in order to record and evaluate the evolution of fibropapilloma in New Caledonia. Necropsy findings confirmed that stranded individuals were in poor health condition but did not succumb directly from fibropapilloma virus. Tumors were generally located on the eyes and on the lower shell disrupting locomotion and vision thus enabling their capacity to feed (Herbst [4]; Flint et al. [17]).

40% of the data come from photo-identification from the volunteer group "Fantastic Grandmothers" which allowed monitoring of progression of tumors from sequential photos (Table 1). Among them, seven individuals were seen regularly. Six were observed with tumor growth in only a few months (Figures 2, 3 and 4). One individual "Yana" (Table 1) was seen for the first time in September 2020 and recorded with tumor growth four years later. However, tumor regression was observed in 2025 (Figure 4). This is the first case of tumor regression recorded on an individual in New Caledonia. Spontaneous lesion regression has also been observed in Brazil (Machado Guimarães et al. [44]), USA (Page-Karjian et al. [45]), Australia and Hawaii (Aguirre and Lutz [46,47]).

Tumor scoring on the individuals reported in this study are mostly at the first stage of infection (14 individuals) and only one individual was recorded with a stage 3 (Table 1). This could indicate that the lesions are recent and that the conditions for the growth of these lesions are currently favorable in the waters of New Caledonia with a strong number of reports in the locality of Nouméa. Urbanization works are being done in the area since 2022 with a peak of activity at the end of 2023 and 2024 and could impact water turbidity. These external parameters could explain the higher number of lesions being recorded and the possible regression happening since the activity has settled in the area. Tumor scoring and precise recording of all individuals presenting fibropapilloma lesions are essential in the future in order to assess the evolution of this disease in New Caledonia.

Table 2: Chart of turtles with fibropapilloma reported in New Caledonia by year.


| Year | Number |
|----------|---------|
| 2011 | 4% (1) |
| 2013 | 4% (1) |
| 2014 | 8% (2) |
| 2016 | 8% (2) |
| 2018 | 4% (1) |
| 2019 | 8% (2) |
| 2021 | 8% (2) |
| 2022 | 16% (4) |
| 2024 | 32% (8) |
| Mid-2025 | 8% (2) |

**Figure 2:** Evolution of tumors on the left eye on "Hope":

a) no tumors in December 2023

b) and c) growth of tumors in July 2024


Photographs by: a) Aline Guémas, b) Valérie Brouns, c) Monique Mazière

**Figure 3:** Evolution of tumors on the left fin on "Hope":

d) no tumors in December 2023

e) growth of tumors in July 2024

Photographs by: d) Aline Guémas, e) Monique Mazière

**Figure 4:** Regression of tumors on a green turtle "Yana":

f) no tumors in September 2020

g) tumors in October 2024

h) resorbed tumors in June 2025 <https://www.inaturalist.org/observations/292566691>

Photographs by: f) Aline Guémas, g) Monique Mazière, h) Valérie Brouns

Conclusion

This study highlights the evolution over the last fourteen years of the fibropapilloma virus in New Caledonia. Among all twenty-five reports, only *Chelonia mydas* were impacted and in particular juveniles. The majority of the fibropapilloma reports were recorded from stranding or monitored by photo-identification and located in South province of New Caledonia. The first case of regression is also highlighted in this study. The disease needs to be closely monitored in the future with specific campaigns dedicated to this issue.

Declarations

Acknowledgment: Deepest thanks to the team of scientists, local governments employees in charge of biodiversity (especially conservation officers), members and volunteers of the Aquarium des Lagons, park rangers, all volunteers from the stranding network and the Fantastic Grandmothers who gave incredible work with the photo-identification program and to Jérôme Villemain who contributed to the development of the map. Special thanks to all veterinarians who helped with the rehabilitation of turtles and in memory of our dear co-worker Jean-Christophe Vivier, who passed away before the first stages of publication. We want to thank Thierry Work for his contribution to the writing of the paper and for the data analyses he conducted that helped through this study. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Author's contributions: Roxane Tran performed analysis and wrote the paper. Claire Goiran contributed to the writing and data analyses. Jean-Jérôme Cassan and Olivier Chateau contributed to the writing and to the data collection. Frédéric Avril, Izabella Lach, Agathe Binois and Laurent Fabre contributed to the data collection. Tyffen Read contributed to the writing and the research idea, collected the data and performed analysis.

Funding: No funding for this study.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Whilde J, Mashkour N, Koda SA (2024) Global overview of sea turtle fibropapillomatosis tumors: a survey of expert opinions and trends.
- Smith G, Coates C (1938) Fibro-epithelial growths of the skin in large marine turtles, *Chelonia mydas* (Linnaeus). *Zoologica* 23: 93-98.
- Page-Karjian A (2019) Fibropapillomatosis in Marine Turtles. *Fowler's zoo and wild animal medicine current therapy* 9: 398-403.
- Herbst LH (1994) Fibropapillomatosis of marine turtles. *Annual Review of Fish Diseases* 4:389-425.
- Jones K, Ariel E, Burgess G, Read M (2016) A review of fibropapillomatosis in Green turtles (*Chelonia mydas*). *The Veterinary Journal* 212: 48-57.
- Whitlock PA, Hyatt E, Belskis LC (2025) Proceedings of the 42nd Annual Symposium on Sea Turtle Biology and Conservation 24-29 March, 2024 Pattaya, Thailand Proceedings of the Forty-Second Annual Symposium on Sea Turtle Biology and Conservation. Prevalence, distribution and etiology of fibropapillomatosis in immature green turtles (*Chelonia Mydas*) of the West Indies 19.
- Herbst LH, Klein PA (1995) Green turtle fibropapillomatosis: challenges to assessing the role of environmental cofactors. *Environ Health Perspect* 103: 27-30.
- Lu Y, Yu Q, Zamzow JP (2000) Detection of Green Turtle Herpesviral Sequence in Saddleback Wrasse *Thalassoma duperrey*: A Possible Mode of Transmission of Green Turtle Fibropapilloma. *Journal of Aquatic Animal Health* 12: 58-63.
- Greenblatt RJ, Work TM, Balazs GH (2004) The *Ozobranchus leech* is a candidate mechanical vector for the fibropapilloma-associated turtle herpesvirus found latently infecting skin tumors on Hawaiian green turtles (*Chelonia mydas*). *Virology* 321: 101-110.
- Work TM, Dagenais J, Balazs GH (2015) Dynamics of Virus Shedding and In Situ Confirmation of Chelonid Herpesvirus 5 in Hawaiian Green Turtles with Fibropapillomatosis. *Vet Pathol* 52: 1195-1201.
- Dupont SM, Bustamante P, Duffy DJ (2024) Future research avenues for the study of fibropapillomatosis in sea turtles. *Front Ecol Evol* vol. 12.
- Greenblatt RJ, Quackenbush SL, Casey RN (2005) Genomic Variation of the Fibropapilloma-Associated Marine Turtle Herpesvirus across Seven Geographic Areas and Three Host Species. *J Virol* 79: 1125-1132.
- Adnyana W, Ladds P, Blair D (1997) Observations of fibropapillomatosis in green turtles (*Chelonia mydas*) in Indonesia. *Aust Veterinary J* 75: 737-742.
- Lucero R, Avenido P, Parcasio S (2012) Fibropapillomatosis: A case in marine turtle reared in captivity in the Southern Philippines. *Asian Biologging Science* 65.
- Li T-H, Hsu W-L, Lan Y-C (2017) Identification of Chelonid herpesvirus 5 (ChHV5) in endangered green turtles (*Chelonia mydas*) with fibropapillomatosis in Asia. *bms* 93: 1011-1022.
- Robben DM, Palaniappan P, Loganathan AL, Subbiah VK (2023) Increased Prevalence and New Evidence of Multi-Species Chelonid Herpesvirus 5 (ChHV5) Infection in the Sea Turtles of Mabul Island, Borneo. *Animals* 13: 290.
- Flint M, Limpus CJ, Patterson-Kane JC (2010) Corneal Fibropapillomatosis in Green Sea Turtles (*Chelonia mydas*) in Australia. *Journal of Comparative Pathology* 142: 341-346.
- Ariel E, Nainu F, Jones K, K Juntunen, I Bell, et al. (2017) Phylogenetic Variation of Chelonid Alphaherpesvirus 5 (ChHV5) in Populations of Green Turtles *Chelonia mydas* along the Queensland Coast, Australia. *Journal of Aquatic Animal Health* 29: 150-157.
- Jones K, Limpus CJ, Brodie J (2022) Spatial distribution of fibropapillomatosis in green turtles along the Queensland coast and an investigation into the influence of water quality on prevalence. *Conserv Sci and Prac* 4(8).
- Coudray J (1976) Recherches sur le Néogène et le Quaternaire marins de la Nouvelle-Calédonie: contribution de l'étude sédimentologique à la connaissance de l'histoire géologique post-éocène de la Nouvelle-Calédonie. Etude des algues calcaires de la série quaternaire traversée par le sondage Ténie (Côte Sud-Ouest de la Nouvelle-Calédonie).
- Ouillon S, Douillet P, Fichez R, Panché J-Y (2005) Enhancement of regional variations in salinity and temperature in a coral reef lagoon, New Caledonia. *Comptes Rendus Géoscience* 337: 1509-1517.

22. Rougerie F (1995) Nature et fonctionnement des atolls des Tuamotu (Polynésie Française). *Oceanologica Acta* 18: 61-78.
23. Xun S (2018) The mineral industry of New Caledonia. *Minerals Yearbook: Area Reports: International Review 2014 Asia and the Pacific* 3:15.
24. Bird EC, Dubois JP, Iltis JA (1984) The impacts of opencast mining on the rivers and coasts of New Caledonia. Tokyo: United Nations University 505.
25. Ouillon S, Douillet P, Lefebvre JP (2010) Circulation and suspended sediment transport in a coral reef lagoon: The south-west lagoon of New Caledonia. *Marine Pollution Bulletin* 61: 269-296.
26. Read TC, Fitz Simmons NN, Wantiez L (2015) Mixed stock analysis of a resident green turtle, *Chelonia mydas*, population in New Caledonia links rookeries in the South Pacific. *Wildl Res* 42: 488.
27. Fretey J, Read TC, Carron L (2023) From terra incognita to hotspot: the largest South Pacific green turtle nesting population in the forgotten reefs of New Caledonia. *Oryx* 57: 626-636.
28. Barbier M, Lafage D, Bourgogne H (2023) Assessment of the nesting population demography of loggerhead turtles (*Caretta caretta*) in La Roche Percée: First long-term monitoring in New Caledonia. *Aquatic Conservation* 33: 579-591.
29. Bourgogne H, Oremus M, Mangeas M (2024) Innovative monitoring scheme adapted to remote, scattered nesting aggregation reveals a major loggerhead turtle rookery in New Caledonia, South Pacific.
30. Limpus C, Miller J, Paramenter C (1992) Migration of green (*Chelonia mydas*) and loggerhead (*Caretta caretta*) turtles to and from eastern Australian rookeries. *Wildl Res* 19: 347 -357.
31. Read TC, Wantiez L, Werry JM (2014) Migrations of Green Turtles (*Chelonia mydas*) between Nesting and Foraging Grounds across the Coral Sea. *PLoS ONE* 9: e100083.
32. Cheng IJ (2000) post-nesting migrations of green turtles (*Chelonia mydas*) at Wan-An Island, Penghu Archipelago, Taiwan. *Marine Biology* 137 :747-754.
33. Read T, Wantiez L, Werry JM (2020) Where are the Adults? First Results of a Study on *C. Mydas* Foraging in New Caledonia. *OFOAJ* 11.
34. Work T, Balazs G (2013) Tumors in sea turtles: The insidious menace of fibropapillomatosis. *The Wildlife Professional* 44-47.
35. Limpus C, Couper P, Read M (1994) The green turtle, *Chelonia mydas*, in Queensland: population structure in a warm temperature feeding area. *Memoirs of the Queensland Museum, Brisbane* 35: 139-154.
36. Limpus C, Reed P, Grigg G (1985) The green turtle, *Chelonia mydas*, in Queensland: population structure in a coral reef feeding area.
37. Read T, Farman R, Vivier J-C (2023) Twenty Years of Sea Turtle Strandings in New Caledonia. *Zoological Studies*.
38. Goiran C, Shine R (2019) Grandmothers and deadly snakes: an unusual project in "citizen science." *Ecosphere* 10.
39. Work TM, Balazs GH (1999) Relating Tumor Score to Hematology in Green Turtles with Fibropapillomatosis in Hawaii. *Journal of Wildlife Diseases* 35: 804-807.
40. Balazs G (1991) Current status of fibropapillomatosis in the Hawaiian green turtle, *Chelonia mydas*. *Research for marine turtle fibropapilloma* 47-57.
41. Herbst LH, Jacobson ER, Klein PA (1999) Comparative Pathology and Pathogenesis of Spontaneous and Experimentally Induced Fibropapillomas of Green Turtles (*Chelonia mydas*). *Vet Pathol* 36: 551-564.
42. Read T, Jean C (2021) Using social media and photo-identification for sea turtles of New Caledonia. *Marine Turtle Newsletter* 162: 25-29.
43. EneA, SuM, LemaireS (2005) Distribution of chelonid fibropapillomatosis-associated herpesvirus variants in Florida: molecular genetic evidence for infection of turtles following recruitment to neritic developmental habitats. *Journal of Wildlife Diseases* 41: 489-497.
44. Machado Guimarães S, Mas Gitirana H, Vidal Wanderley A (2013) Evidence of regression of fibropapillomas in juvenile green turtles *Chelonia mydas* caught in Niterói, southeast Brazil. *Dis Aquat Org* 102: 243-247.
45. Page-Karjian A, Norton TM, Krimer P (2014) Factors influencing survivorship of rehabilitating green sea turtles (*Chelonia mydas*) with fibropapillomatosis. *Journal of Zoo and Wildlife Medicine* 45: 507-519.
46. Aguirre AA, Lutz Peter L (2004) Marine Turtles as Sentinels of Ecosystem Health: Is Fibropapillomatosis an Indicator? *EcoHealth* 1(3):275-283.
47. Jones SM, Caspi I, Lucas C (2021) Fibropapillomatosis infection in a population of green turtles at Watamu Bay, Kenya. *West Ind Oc J Mar Sci* 20: 111-123.