

30–32 °C, and both species spent most time in water shallower than 30 m. With 200 satellite tags deployed and a decade of building local research collaborations, we identified that the Mariana Archipelago functions primarily as developmental habitat for range-restricted juvenile green and hawksbill turtles that reside in shallow coastal waters for years. EPUE maps and telemetry together identify discrete nearshore habitats that are likely sensitive to local stressors and therefore high priorities for marine spatial planning and management. This project was a monumental effort which yielded the greatest understanding of foraging behavior for these two endangered (US ESA) species in the Pacific Islands Region.

ENVIRONMENTAL AND HUMAN INFLUENCES ON GREEN SEA TURTLE HABITAT USE IN THE RARE BRACKISH RIVER, OKINAWA, JAPAN


Yuka Matsukawa¹, Jeffrey Jolly², Timothy Ravasi², James Reimer¹

¹University of the Ryukyus, Japan

²Okinawa Institute of Science and Technology (OIST), Japan

Shiokawa River, located on the Motobu Peninsula of Okinawa, Japan, is one of only two known salt water rivers globally, and it has been designated as a Natural Monument of Japan. Shiokawa is scientifically classified as an anchialine spring. This unique brackish ecosystem, with seawater gushing out from underground springs, has long been known by local people as a site where green sea turtles (*Chelonia mydas*) ascend the river. However, scientific understanding of the environmental ecology in such habitats remains extremely limited. Investigating sea turtle behavior in this unique salt river is critical for species conservation and provides insight into ecological interactions in rare brackish habitats worldwide. From September 2024 to August 2025, we conducted observations of sea turtle sightings and environmental information along the Shiokawa River, from sunrise to sunset across different tidal stages. Water temperature in Shiokawa remained stable throughout the year. Sea turtle sightings were concentrated in the early morning and late afternoon, suggesting that habitat use may be influenced not by temperature fluctuations, but by tidal accessibility and reduced human disturbance. These patterns suggest that Shiokawa's role as a sea turtle habitat is shaped more by its unique physical features and patterns of human use. The consistent water temperature quiet conditions during early morning and late afternoon hours, may provide favorable conditions for resting or foraging. Based on individually identified sea turtles, at least seven individuals ascended the river during the study period. Furthermore, photographic records confirmed that at least one individual sea turtle has used the river mouth repeatedly since 2023, indicating strong site fidelity. This study provides the first record of sea turtle behavior in a salt river, and underscores the importance of integrating community-based monitoring to support conservation of these exceptional habitats. These findings demonstrate that even small and rare salt water rivers can provide essential ecological resources for sea turtle. By combining systematic observation with community engagement, this study contributes not only to species conservation but also to a broader understanding of globally rare brackish river ecosystems. Continued monitoring and sustained collaboration with residents are needed to understand long-term habitat use and to support conservation efforts.

PROCEEDINGS OF THE FORTY-FOURTH SYMPOSIUM ON SEA TURTLE BIOLOGY AND CONSERVATION



February 28 – March 6, 2026
Kailua-Kona, Hawai'i, USA

Compiled by:
Paul A. Whittock, Lisa C. Belskis, & Joseph B. Pfaller

June 2026